arXiv:1609.02988v1 [math.AG] 10 Sep 2016

GROUP SCHEMES OF SQUARE FREE ORDER

V. KUMAR MURTY, YING ZONG

Let S be a scheme and let $f : G \to S$ be a finite flat finitely presented S-group scheme. The *order*, or *rank*, of G, namely, the rank of the locally free $\mathcal{O}_S$-module of finite type $f_*\mathcal{O}_G$, is a locally constant function on S. We say that G is of *prime order* (resp. *square free order*) if its order at each point of S is 1 or a prime (resp. non-divisible by prime squares).

Theorem. — *Let S be a scheme and let G be a finite flat finitely presented S-group scheme of square free order. Then there exists a Hochschild extension of S-group schemes*

$$1 \to G' \to G \to G'' \to 1,$$

split by a finite étale surjective base change, with G'' finite étale over S, and G' a direct sum, as G''-modules, of commutative finite flat finitely presented S-group schemes of prime order. If

$$1 \to G'_\iota \to G \to G''_\iota \to 1,$$

$\iota = 1, 2$, are two such extensions, there is a third

$$1 \to G'_3 \to G \to G''_3 \to 1$$

with $G'_3 = G'_1 \cap G'_2$.

Lemma. — *Let S be a scheme and let G be a finite flat finitely presented S-group scheme of square free order. Then :*

a) If G is of prime order, it is commutative.

b) If G is commutative and of order n, it is in a unique way a direct sum, indexed by the factors $p \in \Gamma(S, \mathbf{Z})$ in the prime factorization of n, of finite flat finitely presented S-group schemes of prime order p. This decomposition commutes with base change and is invariant by all S-group automorphisms of G.

c) Each extension of S-group schemes

$$1 \to G' \to G \to G'' \to 1,$$

with G'' S-finite étale and G' commutative finite flat of finite presentation over S, is Hochschild and split by a finite étale surjective base change.

d) If, for $\iota = 1, 2$, either

$$1 \to G'_\iota \to G \to G''_\iota \to 1$$

writes G as extension of a finite étale S-group scheme by a finite flat finitely presented S-group scheme, then there is a third such

$$1 \to G' \to G \to G'' \to 1$$

with $G' = G'_1 \cap G'_2$.

e) If S is of residue characteristic zero, G is finite étale over S.

f) If G is finite étale over S, then, for each $d \in \Gamma(S, \mathbf{Z})$, the set U of points s of S such that $G(\overline{s})$ has a unique subgroup of order $d(s)$, where $\overline{s}$ is the spectrum of an algebraic closure of $k(s)$, is open and closed in S. Over U, $G \times_S U$ has a unique normal finite étale sub-U-group scheme of order d.

g) If S is the spectrum of an algebraically closed field of characteristic $p > 0$, the identity component of G, if non-trivial, is μ_p or α_p.

h) For every point s of S, if $i_G(s)$ denotes the order of the identity component of $G \times_S s$, the function $s \mapsto i_G(s)$, called the infinitesimal rank of G, is locally constructible upper semi-continuous.

i) The set S_1 of points s of S such that $i_G(s)$ is 1 is a retro-compact open subset of S; it consists exactly of all those points of S above which G is S-étale.

j) Let p be a prime number. The set S_p of points s of S such that $i_G(s)$ divides p is a retro-compact open subset of S. The union V_p of $S_p - S_1$ and the points s of S such that $G(\overline{s})$ has a unique p-Sylow subgroup of order p, where $\overline{s}$ is the spectrum of an algebraic closure of $k(s)$, is open and closed in S. On V_p, $G \times_S V_p$ has a unique normal finite flat finitely presented sub-V_p-group scheme G_p of order p.

Proof. *a*) This is [1] §1 Theorem 1.

b) By [1] §1 Theorem, as an abelian sheaf on S for the *fppf* topology, G is of n-torsion, namely, a module over the locally constant sheaf of rings

$$\mathbf{Z}/n\mathbf{Z} = \prod_p \mathbf{Z}/p\mathbf{Z},$$

and in particular G is the direct sum of its p-torsion sub-modules G_p, where $p \in \Gamma(S, \mathbf{Z})$ are the factors in the prime factorization of n. As the kernel of $p.\mathrm{Id}_G$, G_p is a finitely presented closed sub-S-group scheme of G. As also the *fppf* image of $(n/p).\mathrm{Id}_G$, G_p is S-flat (EGA IV 11.3.11). The order n_p of G_p is p, for

$$n = \prod p \mid \prod n_p = n.$$

This decomposition of G is clearly invariant by all its S-group automorphisms and commutes with base change. The uniqueness follows as every finite flat finitely presented S-group scheme of prime order p is of p-torsion.

c) One may assume G'' of constant order n''. The extension

$$1 \to G' \to G \to G'' \to 1$$

is Hochschild, namely, the sequence

$$1 \to G'(T) \to G(T) \to G''(T) \to 1$$

is exact for every S-scheme T. Indeed, for all $x \in G''(T)$, if

$$d : G''(T) \to H^1((\mathrm{Sch}/T)_{\mathrm{fppf}}, G')$$

denotes the coboundary homomorphism, one has $d(x) = 0$, as

$$n''.d(x) = d(x^{n''}) = d(1) = 0$$

and $n''.\mathrm{Id}_{G'} : G' \to G'$ is an isomorphism. Next, by a finite étale surjective base change, one may assume that the finite étale S-group scheme G'' is locally constant and then constant of value $|G''|$. It suffices to show that the exact sequence

$$1 \to G'(S) \to G(S) \to G''(S) \to 1$$

is split by $|G''| \to G''(S)$. This follows as $H^2(B|G''|, G'(S)) = 0$.

d) The intersection $G' = G'_1 \cap G'_2$ is a closed finitely presented normal sub-S-group scheme of G. It remains to show that G' is flat over S and that the quotient $G'' = G/G'$, which is then representable by a finite flat finitely presented S-group scheme, is étale over S.

The assertion being local on S, one may assume S affine, then by a "passage à la limite", noetherian and local, then by completion along its closed point, complete, and finally by reduction to each of its closed sub-schemes of finite lengths, artin local. Let the closed point of S be s and let G^o_s be the identity component of $G_s = G \times_S s$. Consider the canonical exact sequence

$$1 \to G^o_s \to G_s \to G_s/G^o_s \to 1.$$

As the base change by $s \to S$ induces an equivalence from the category of finite étale S-group schemes to the category of finite étale s-group schemes (EGA IV 18.1.2), there is up to unique isomorphisms a unique finite étale S-group scheme Q with $Q \times_S s = G_s/G^o_s$. By EGA IV 18.1.3, there is a unique S-group homomorphism $G \to Q$ which specializes to

the projection $G_s \to G_s/G_s^o$. This morphism $G \to Q$ is faithfully flat (EGA IV 11.3.11) and provides an exact sequence of S-group schemes

$$1 \to P \to G \to Q \to 1$$

with P finite flat over S, $P \times_S s = G_s^o$. For $\iota = 1, 2$, either composition

$$P \to G \to G_\iota''$$

is by EGA IV 18.1.3 trivial, for it is when specialized to s. So $P \subset G_1' \cap G_2' = G'$. Replacing G by G/P, G_ι' by G_ι'/P and G' by G'/P, one can assume $G = Q$ finite étale over S. But now all of G_1', G_2', G', G'' are finite étale over S. This completes the proof.

e) One can (EGA IV 17.6.2) assume that S is the spectrum of a field of characteristic zero and apply SGA 3 VII B 3.3.1.

f) The question being local on S, one can assume S affine and by a "passage à la limite" noetherian and connected. If U is not empty, let $\overline{s}$ be a geometric point of S with image s in U and let

$$\rho : \pi_1(S, \overline{s}) \to \mathrm{Aut}(G(\overline{s}))$$

be the monodromy representation corresponding to the finite étale S-group scheme G. The unique subgroup of $G(\overline{s})$ of order $d(s)$ is characteristic in $G(\overline{s})$ and in particular invariant by ρ. So G has a unique normal finite étale sub-S-group scheme of order d, and $U = S$.

g) This follows by SGA 3 XVII 4.2.1.

h) The order of the quotient of $G \times_S s$ by its identity component, the *separable rank* of $G \times_S s$, is a locally constructible lower semi-continuous function of $s \in S$ (EGA IV 15.5.1).

i) This follows by *h*) and EGA IV 17.6.2.

j) That S_p is a retro-compact open subset of S follows by *g*)+*h*).

— *The set $U_p := V_p \cap S_p$ is open and closed in S_p, and $G \times_S U_p$ has a unique normal finite flat finitely presented sub-U_p-group scheme P of order p. The quotient $(G \times_S U_p)/P$ is finite étale over U_p* :

Restricting to S_p, suppose $S = S_p$, namely, that i_G takes values in $\{1, p\}$. Let $U : (\mathrm{Sch}/S)^o \to (\mathrm{Sets})$ be the following sub-functor of the final functor :

For an S-scheme S', $U(S') = \{\emptyset\}$, if $G \times_S S'$ is the extension of a finite étale S'-group scheme by a finite flat finitely presented S'-group scheme of order p, and $U(S') = \emptyset$, otherwise.

It suffices to show that $U \to S$ is representable by an open and closed immersion.

The functor U is a sheaf on (Sch/S) *for the étale topology* :

Namely, $U(T) = U(T')$ for every étale surjective morphism $T' \to T$ of S-schemes. Indeed, if $U(T')$ is not empty so that there is an extension of T'-group schemes

$$1 \to P' \to G \times_S T' \to Q' \to 1$$

with Q' T'-finite étale and P' finite flat of finite presentation over T' of order p, it follows by d) that $p_1^* P' = p_2^* P'$ in $G \times_S T''$, where p_1, p_2 are the two projections of $T'' = T' \times_T T'$ onto T'. On $P' \hookrightarrow G \times_S T'$, there is therefore a canonical descent datum relative to $T' \to T$, and so $U(T)$ is not empty.

The functor U is formally étale :

Namely, $U(T) = U(T')$ for every nilpotent S-immersion $T' \hookrightarrow T$. This is immediate from EGA IV 18.1.2+18.1.3 by a similar argument as in d).

The functor U verifies the valuative criterion of properness :

As U is a sheaf on (Sch/S) for the étale topology, the question is local on S. It suffices to assume S affine and by a "passage à la limite" noetherian. Then given an S-scheme T which is the spectra of a discrete valuation ring and which has generic point t, one has $U(T) = U(t)$. Indeed, if $U(t)$ is not empty and so $G \times_S t$ is the extension of a finite étale t-group scheme Q_t by a finite t-group scheme P_t of order p, then $Q = G_T/P$ is finite étale over T, where P is the closed image of P_t in $G_T = G \times_S T$. For, the infinitesimal rank of G, hence that of Q as well, takes values in $\{1, p\}$, and so $i_Q(T) = \{1\}$, as Q is of order prime to p.

Now it is clear that U is representable by an open and closed subscheme of S with underlying set U_p.

— *The set $U'_p := V_p - (S_p - S_1)$ is open and closed in $S - (S_p - S_1)$, and $G \times_S U'_p$ has a unique normal finite étale sub-U'_p-group scheme of order p* :

Restricting to each S_q, where q is a prime distinct from p, one may assume $S = S_q$. And by f), it suffices to assume $S = U_q$. Consider the canonical exact sequence

$$1 \to Q \to G \to G/Q \to 1$$

where Q is the unique normal finite flat finitely presented sub-S-group scheme of G of order q. By f) applied to the prime-to-q finite étale quotient G/Q, and replacing S by an open and closed sub-scheme, one may assume G/Q has a unique normal finite étale sub-S-group scheme $\overline{P}$ of order p. Let $\overline{G}$ be the pre-image of $\overline{P}$ in G. Note that replacing G by $\overline{G}$ one may assume $P := G/Q$ of order p. According to c), by a finite étale surjective base change, one may assume that $G = QP$ is

a semi-direct product and that P is constant of generator g. Now, by conjugation, P acts on Q. Let ζ be the image of g in $\mathrm{Aut}_S(Q)$ and consider the following sub-functor of the final functor :

For an S-scheme S', $V(S') = \{\emptyset\}$, if $\zeta \times_S S' = \mathrm{id}_Q \times_S S'$, and $V(S') = \emptyset$, otherwise.

This sub-functor $V \to S$ is representable by a closed immersion (SGA 3 VIII 6.4). And V has underlying set U'_p, and $G \times_S V$ is commutative with its p-torsion being the unique normal finite étale sub-V-group scheme of order p.

On S_1, $V \times_S S_1 \to S_1$ is an open and closed immersion. For, $Q \times_S S_1$, hence $\underline{\mathrm{Aut}}_S(Q) \times_S S_1$ as well, is finite étale over S_1.

To show that $V \to S$ is an open immersion at each point s of $U'_p - S_1$, it suffices by strictly localizing S at s to assume S strictly local. By [1] §2 Theorem 2, Q corresponds to a triple (L, c, d) which consists of an invertible $\mathcal{O}_S$-module L and of $\mathcal{O}_S$-linear homomorphisms $c : L \to L^{\otimes q}$, $d : L^{\otimes q} \to L$ satisfying $d \circ c = w.\mathrm{Id}_L$ for a certain element $w \in \Gamma(S, \mathcal{O}_S)$. This correspondence identifies the S-group automorphisms of Q with the units $u \in \Gamma(S, \mathcal{O}_S)^\times$ such that $uc = u^q c$, $ud = u^q d$. In particular, ζ is identified to an element of $\mu_p(S)$, and the relation "$\zeta = \mathrm{Id}_Q$" is an open and closed relation on S, as μ_{pS} is finite étale over S.

□

Proof of the theorem —

The last assertion follows by Lemma, d). By Lemma, c)+b), it only remains to write G as an extension

$$1 \to G' \to G \to G'' \to 1$$

with G'' S-finite étale and G' commutative finite flat of finite presentation over S.

One may assume G of constant order n. The order $i_G(s)$ of the identity component of each fiber $G \times_S s$ divides n and equals either 1 or the characteristic of $k(s)$. If $E := i_G(S)$ consists only of 1, G is S-finite étale (Lemma, i)). Otherwise, for each prime p in E, there is (Lemma, j)) an open and closed sub-scheme V_p of S such that :

— *On $S - V_p$, i_G does not take value p.*

— *On V_p, $G \times_S V_p$ has a unique normal finite flat finitely presented sub-V_p-group scheme G_p of order p.*

By induction on the cardinality of $E - \{1\} = \{p_1, p_2, \cdots\}$, suppose that the claim holds for the restriction of G to

$$S - (V_{p_1} \cap V_{p_2} \cap \cdots) = (S - V_{p_1}) \coprod (V_{p_1} - V_{p_2}) \coprod (V_{p_1} \cap V_{p_2} - V_{p_3}) \coprod \cdots$$

Then, restricting to $V = V_{p_1} \cap V_{p_2} \cap \cdots$, one may assume $V_p = S$ for every prime p in E. Put $G' = G_{p_1} \times_S G_{p_2} \times_S \cdots$. Notice that the S-morphism

$$G' \to G,\ (g_1, g_2, \cdots) \mapsto g_1.g_2 \cdots$$

identifies G' as a normal sub-S-group scheme of G with $G'' = G/G'$ finite étale over S.

References

1. F. Oort, J. Tate. Group schemes of prime order. *Ann. scient. Éc. Norm. Sup.* 4^e série, t. 3, 1970.

Department of Mathematics, University of Toronto
E-mail address: murty@math.utoronto.ca, zongying@math.utoronto.ca